



**IN THE CLAIMS:**

Please amend the claims, as follows:


Claim1(currently amended): A method of ~~manufacturing~~ preparation of 7-ethyl-10-[4-(1-piperidino)-1-piperidino]-carbonyloxy-camptothecin of formula I



~~characterized in that~~ wherein 7-ethyl-10-hydroxycamptothecin of formula II



is subjected to a condensation reaction with 1-chlorocarbonyl-4-piperidinopiperidine hydrochloride of formula III



in a polar aprotic solvent, e.g. in acetonitrile, in the presence of 4-dimethylaminopyridine.

Claim 2 (currently amended): The method according to claim 1, characterized in that wherein 1-chlorocarbonyl-4-piperidinopiperidine hydrochloride is employed in an amount of 1.3 to 3 mol, preferably in an amount of 1.6 to 1.9 mol, per 1 mol of 7-ethyl-10-hydroxycamptothecin.

Claim 3 (currently amended): The method according to claim 1, characterized in that wherein 4-dimethylaminopyridine is employed in an amount of 1.5 to 4 mol, preferably in an amount of 1.8 to 2.2 mol, per 1 mol of 7-ethyl-10-hydroxycamptothecin.

Claim 4 (currently amended): The method according to claim 1, characterized in that wherein the polar aprotic solvent is employed in an amount of 400 to 600 mol, preferably in an amount of 430 to 460 mol, per 1 mol of 7-ethyl-10-hydroxycamptothecin.

Claim 5 (currently amended): The method according to claim 1, characterized in that wherein the condensation reaction is carried out at a temperature of 70 to 80 °C, preferably at a temperature of 73 to 77 °C.